

p-Wave Resonant Bose Gas: A Finite-Momentum Spinor Superfluid

Leo Radzihovsky and Sungsoo Choi

Department of Physics, University of Colorado, Boulder, CO 80309

(Dated: August 31, 2009)

We show that a degenerate gas of two-species bosonic atoms interacting through a *p*-wave Feshbach resonance (as realized in, e.g., a ^{85}Rb - ^{87}Rb mixture) exhibits a *finite-momentum* atomic-molecular superfluid (AMSF), sandwiched by a molecular *p*-wave (orbital spinor) superfluid and by an *s*-wave atomic superfluid at large negative and positive detunings, respectively. The magnetic field can be used to tune the modulation wave vector of the AMSF state, as well as to drive quantum phase transitions in this rich system.

A Feshbach resonance (FR) is an exceptionally fruitful experimental “knob” that allows exquisite tunability of interactions in degenerate atomic gases. This has led to realizations and studies of Bose-Einstein condensation (BEC)-BCS crossover of fermion-paired *s*-wave superfluidity [1, 2, 3, 4]. The bosonic counterparts have also been extensively explored, and in fact, in the *s*-wave FR case, e.g., in ^{85}Rb [5], predate recent fermionic developments. As was recently emphasized [6, 7, 8], in contrast to their fermionic analogs that undergo a smooth BEC-BCS crossover, resonant bosonic gases are predicted to exhibit magnetic field and/or temperature driven sharp phase transitions between distinct molecular and atomic superfluid phases.

Motivated by these successes, recent attention has focused on a realization of an even richer *p*-wave paired fermionic superfluidity [9, 10, 11, 12], utilizing *p*-wave FR in ^{40}K and ^6Li . Laboratory production of *p*-wave Feshbach molecules [13, 14] showed considerable promise toward this goal; however, reaching molecular degeneracy has been plagued with short molecular lifetimes.

In another important development, experiments on a ^{85}Rb - ^{87}Rb mixture have demonstrated a *p*-wave FR at $B = 257.8\text{G}$ between these two bosonic isotopes [15]. Although consequences of this two-body *p*-wave resonance on the degenerate state of such a gas mixture has not been further explored experimentally, it provides the main motivation for our work. In this Letter, we report on our study of a two-species degenerate Bose gas with a *p*-wave Feshbach resonant interspecies interaction.

As summarized by the phase diagram in Fig. 1, we find (within a mean-field treatment that we expect to largely survive fluctuations) that in addition to the normal (N, i.e., nonsuperfluid) phase, the *p*-wave Feshbach resonant two-component *balanced* Bose gas (e.g., equal mixture of ^{85}Rb and ^{87}Rb atoms) exhibits three classes of superfluid phases: atomic (ASF), molecular (MSF), and atomic-molecular (AMSF) condensates. Our most interesting finding is that the AMSF, sandwiched between (large positive detuning) ASF and (large negative detuning) MSF phases is necessarily a *finite-momentum* Q spinor superfluid, with a characteristic wave vector (with

$\hbar = 1$)

$$Q = \alpha m \sqrt{n_m} \sim \sqrt{\gamma_p \ell n_m} \lesssim \sqrt{\gamma_p} / \ell, \quad (1)$$

tunable with a magnetic field (via FR detuning, ν that primarily enters through the molecular condensate density $n_m(\nu)$), with α , m , ℓ , and γ_p , respectively, the FR coupling, atomic mass, atom spacing, and a dimensionless measure of FR width[11].




FIG. 1: (color online). Schematic temperature-detuning phase diagram for a two-species mixture of bosonic atoms exhibiting atomic (ASF), molecular (MSF), and atomic-molecular (AMSF) phases. In the AMSF *p*-wave, molecular condensate coexists with a finite-momentum Q (see left inset) atomic superfluid. The cartoon in AMSF phase illustrates the mechanism driving finite-momentum condensation.

Within the narrow ($\gamma_p \ll 1$) FR approximation[6, 7, 10, 11], we find that the optimum collinear state is characterized by a single Q , Fulde-Ferrell-like state [16], as opposed to a $+Q$ and $-Q$ Larkin-Ovchinnikov-like state[17] found in imbalanced paired fermionic systems[18, 19, 20]. However, a more detailed study is necessary to ascertain the precise set of Q ’s at which AMSF condensation takes place.

The physical picture behind the finite-momentum AMSF formation is illustrated in Fig. 1. At intermediate detuning where atomic gap closes within the MSF state, *p*-wave molecules decay via FR into a pair of atoms, which (due to the *p*-wave nature of the molecules) are necessarily created at finite and opposite momenta,

$\pm \mathbf{k}$, and therefore at low temperature form a finite-momentum atomic condensate, AMSF. The energetic cost ($\sim k^2/2m$) of a finite momentum atomic condensation is balanced by the lowering of the energy ($\sim \alpha k \sqrt{n_m}$) through FR hybridization between closed-channel p -wave molecule and open-channel pair of atoms that is only possible at finite atomic momentum k , giving Q in Eq. (1).

As we detail below, in addition, we find that each of the three superfluid classes (ASF, MSF, AMSF) in turn consist of distinct phases selected by detuning, temperature, and background s -wave scattering lengths, and distinguished by the nature (ferromagnetic or polar) of the p -wave molecular condensate and/or which combination of the two types of atoms is Bose condensed. We explore the nature of these SF phases and associated phase transitions.

To outline the derivation of these results, we consider a model of a gas mixture of two distinguishable bosonic atoms (e.g., ^{85}Rb , ^{87}Rb) [15], created by field operators $\hat{\psi}_\sigma^\dagger(\mathbf{r}) = (\hat{\psi}_1^\dagger(\mathbf{r}), \hat{\psi}_2^\dagger(\mathbf{r}))$, and interacting through a p -wave Feshbach resonance associated with a tunable closed-channel bound state. The corresponding p -wave ($\ell = 1$) closed channel hetero-molecule (e.g., ^{85}Rb - ^{87}Rb) is created by a vector field operator $\hat{\phi}^\dagger(\mathbf{r}) = (\hat{\phi}_x^\dagger, \hat{\phi}_y^\dagger, \hat{\phi}_z^\dagger)$. This system is governed by a grand-canonical Hamiltonian $H[\hat{\psi}_\sigma, \hat{\phi}] = \int d^3r \mathcal{H}$, with

$$\mathcal{H} = \sum_{\sigma=1,2} \hat{\psi}_\sigma^\dagger \hat{\varepsilon}_\sigma \hat{\psi}_\sigma + \hat{\phi}^\dagger \cdot \hat{\omega} \cdot \hat{\phi} + \mathcal{H}_{bg} \quad (2)$$

$$+ \frac{\alpha}{2} \hat{\phi}^\dagger \cdot [\hat{\psi}_1(-i\nabla)\hat{\psi}_2 - \hat{\psi}_2(-i\nabla)\hat{\psi}_1] + H.c.,$$

where $\hat{\varepsilon}_\sigma = -\frac{1}{2m}\nabla^2 - \mu_\sigma$, $\hat{\omega} = -\frac{1}{4m}\nabla^2 - \mu_m$ with the molecular chemical potential $\mu_m = \mu_1 + \mu_2 - \nu$ adjustable by detuning ν . We have taken atomic masses to be identical (a good approximation for the ^{85}Rb - ^{87}Rb mixture) [21], and focus on the balanced case of $\mu_1 = \mu_2 = \mu$, with μ fixing the total number of ^{85}Rb and ^{87}Rb atoms, whether in the (open-channel) atomic or (closed-channel) molecular form.

The FR interaction encodes a coherent interconversion between a pair of open-channel atoms 1, 2 (in a singlet combination of 1, 2 labels, as required by bosonic statistics) and a closed-channel p -wave molecule, with amplitude α [21]. We have focused on a rotationally invariant FR interaction, with $\hat{\omega}$ and α independent of the molecular component i . This is an approximation to ^{85}Rb - ^{87}Rb mixture, where the p -wave FR around $B = 257.8\text{G}$ is split into a doublet by $\Delta B = 0.6\text{G}$, similarly to the fermionic case of ^{40}K [10, 11, 13, 22]. We leave the more realistic, richer case for future studies [23].

The FR coupling α and detuning ν are fixed experimentally through measurements of the low-energy p -wave scattering amplitude [13, 22] $f_p(k) = \frac{k^2}{-v^{-1} + \frac{1}{2}k_0 k^2 - ik^3}$. The scattering volume, v (tunable via ν) and the characteristic wave vector, k_0 (a p -wave analog of the effective

range, negative for the FR case) [11] define our model parameters in terms of these experimental observables.

The background (nonresonant) interaction density $\mathcal{H}_{bg} = \mathcal{H}_a + \mathcal{H}_m + \mathcal{H}_{am}$ consists of

$$\mathcal{H}_a = \sum_{\sigma=1,2} \frac{\lambda_\sigma}{2} \hat{\psi}_\sigma^\dagger \hat{\psi}_\sigma^2 + \lambda_{12} \hat{\psi}_1^\dagger \hat{\psi}_2^\dagger \hat{\psi}_2 \hat{\psi}_1, \quad (3)$$

$$\mathcal{H}_m = \frac{g_1}{2} (\hat{\phi}^\dagger \cdot \hat{\phi})^2 + \frac{g_2}{2} |\hat{\phi} \cdot \hat{\phi}|^2, \quad (4)$$

$$\mathcal{H}_{am} = \sum_{\sigma=1,2} g_{am} \hat{\psi}_\sigma^\dagger \hat{\phi}^\dagger \cdot \hat{\phi} \hat{\psi}_\sigma, \quad (5)$$

where coupling constants λ_σ , λ_{12} , $g_{1,2}$, g_{am} are related to the corresponding s -wave scattering lengths (a_1 , a_2 , etc.) in a standard way, and thus are fixed experimentally through measurements on the gas in a dilute limit. The miscibility of a two-component atomic gas requires [24] $a_1 a_2 > a_{12}^2$, which may be problematic for the case of ^{85}Rb - ^{87}Rb due to the negative background scattering length of ^{85}Rb .

The molecular interaction couplings g_1 , g_2 (set by the $L = 0$ and $L = 2$ channels of p -wave molecule-molecule scattering) and g_{am} can be derived from a combination of s -wave atom-atom (λ_σ) and p -wave FR (α) interactions [23].

Qualitative features of the phase diagram for the system can be mapped out through a mean-field treatment of the Hamiltonian (2). This amounts to a minimization of the Landau free-energy functional $F[\Psi_\sigma, \Phi]$ of classical fields $\Psi_\sigma(\mathbf{r})$, $\Phi(\mathbf{r})$, corresponding to the coherent state field configurations for the atomic and molecular operators. In the simplest approximation, $F[\Psi_\sigma, \Phi]$ takes the form identical to $H[\hat{\psi}_\sigma, \hat{\phi}]$, with the effective couplings ($\tilde{\mu}_\sigma, \tilde{\mu}_m, \tilde{\lambda}_\sigma, \dots$) that are functions of microscopic parameters ($\mu_\sigma, \nu, \lambda_\sigma, \dots$) appearing in (2).

Minimization of $F[\Psi_\sigma, \Phi]$ is quite straightforward [23]. For large *positive* detuning ν , closed-channel molecules are gapped, and the ground state is a molecular vacuum. Thus, at low temperature $\tilde{\mu}_m < 0$ and $F[\Psi_\sigma, \Phi]$ is minimized by $\Phi = 0$, reducing to $F_a[\Psi_\sigma] = F[\Psi_\sigma, 0] = \int d^3r \left[\sum_{\sigma=1,2} \left(\Psi_\sigma^* \hat{\varepsilon}_\sigma \Psi_\sigma + \frac{\tilde{\lambda}_\sigma}{2} |\Psi_\sigma|^4 \right) + \tilde{\lambda}_{12} |\Psi_1|^2 |\Psi_2|^2 \right]$. This functional is a special ($U(1) \times U(1)$) case of a $O(N) \times O(M)$ model that has been studied extensively [25, 26, 27]. The free-energy is clearly minimized by uniform Ψ_1, Ψ_2 , as this lowers atomic kinetic energy. For $\tilde{\lambda}_1 \tilde{\lambda}_2 > \tilde{\lambda}_{12}^2$ in addition to the normal (nonsuperfluid) state, the system exhibits three ASF phases: (i) ASF₁ with $\Psi_1 \neq 0, \Psi_2 = 0$, (ii) ASF₂ with $\Psi_1 = 0, \Psi_2 \neq 0$, (iii) ASF₁₂ with $\Psi_1 \neq 0, \Psi_2 \neq 0$, separated by continuous phase transitions. For a balanced mixture $\tilde{\mu}_1 = \tilde{\mu}_2$, the system exhibits a direct N-ASF₁₂ transition through a tetracritical point, $\tilde{\mu}_1 = \tilde{\mu}_2 = 0$, that is believed to be in the decoupled universality class [25, 26, 27]. For $\tilde{\lambda}_1 \tilde{\lambda}_2 < \tilde{\lambda}_{12}^2$, the ASF₁₂ phase is absent, and ASF₁ and ASF₂ are separated by a

first-order transition that terminates at a bicritical point[25, 26, 27]. All other transitions (N-ASF₁, N-ASF₂, and ASF_i-ASF₁₂) are in the *XY* universality class, breaking associated *U*(1) symmetries. The phase boundaries and the values of the atomic condensate order parameters can be straightforwardly computed within mean-field theory (MFT)[23], but are modified by fluctuations[26, 27].

Within ASF phases, the spectrum of fluctuations can be straightforwardly computed by a Bogoliubov diagonalization of coupled atomic and molecular excitations, with details depending on which of the three possible ASF phases is studied. In general, there will be one Bogoliubov sound mode per broken atomic *U*(1) symmetry, with one Goldstone mode in ASF₁ and ASF₂ phases and two in ASF₁₂[23].

In the opposite limit of large *negative* detuning, ν open-channel atoms are gapped, and the ground state is an atomic vacuum. Hence, at low temperature $\tilde{\mu} < 0$ and $F[\Psi_\sigma, \Phi]$ is minimized by $\Psi_\sigma = 0$ and a uniform molecular condensate Φ , reducing to $F_m[\Phi]/V = F[0, \Phi]/V = -\tilde{\mu}_m|\Phi|^2 + \frac{g_1}{2}(\Phi^* \cdot \Phi)^2 + \frac{g_2}{2}|\Phi \cdot \Phi|^2$, for (orbital $\ell = 1$) spin-1 molecular bosons with well-studied thermodynamics [28, 29, 30, 31]. In particular, we predict our system to also exhibit polar (MSF_p for $\tilde{g}_2 < 0$) and ferromagnetic (MSF_{fm} for $\tilde{g}_2 > 0$) molecular condensates, respectively, corresponding to $\Phi = \Phi_0 \hat{n} \in [S_2 \times U_N(1)]/Z_2$ and $\Phi = \Phi_0(\hat{n} + i\hat{m})/\sqrt{2} \in SO(3)$ order parameters, with $\hat{n}, \hat{m}, \hat{\ell} \equiv \hat{n} \times \hat{m}$ an orthonormal triad and Φ_0 a complex amplitude, breaking $SO(3) \times U_N(1)$ [32, 33]. The finite *T* N-MSF transitions are in the universality class of a complex *O*(3) model[27].

The Goldstone mode content of these orbital $\ell = 1$ molecular condensate phases is also identical to that of spinor condensates[28], with MSF_p exhibiting three $E_m^{(\text{MSF}_p)}(k) \sim k$ Bogoliubov modes and MSF_{fm} characterized by one $E_{m1}^{(\text{MSF}_{fm})}(k) \sim k$ Bogoliubov and one $E_{m2}^{(\text{MSF}_{fm})}(k) \sim k^2$ ferromagnetic spin-wave modes. An attractive new feature of these orbital molecular condensates is that $\tilde{g}_2(\nu)$ is a tunable function of detuning that can be used to drive a 1st order MSF_p-MSF_{fm} transition[11, 23].

To calculate the spectrum of low-energy excitations inside the MSF (polar and ferromagnetic) phases, we separate molecular field $\hat{\phi} = \Phi + \hat{\varphi}$ into a condensate and small fluctuations about it, obtaining $\mathcal{H}[\hat{\psi}_\sigma, \Phi + \hat{\varphi}] \approx \mathcal{E}_g^{(0)}[\Phi] + \mathcal{H}_a^{(2)} + \mathcal{H}_m^{(2)}$, where $\mathcal{E}_g^{(0)}[\Phi] = \mathcal{H}[0, \Phi]$ is the zeroth-order approximation to the MSF ground state energy,

$$\mathcal{H}_a^{(2)} = \sum_{\sigma=1,2} \hat{\psi}_\sigma^\dagger \tilde{\varepsilon}_\sigma \hat{\psi}_\sigma + \alpha \Phi \cdot \hat{\psi}_1 (-i\nabla) \hat{\psi}_2 + H.c. \quad (6)$$

is the quadratic atomic Hamiltonian density, and $\mathcal{H}_m^{(2)} = \hat{\varphi}_i^\dagger \tilde{\omega}_{ij} \hat{\varphi}_j + \frac{g_1}{2} \Phi_i^* \Phi_j^* \hat{\varphi}_i \hat{\varphi}_j + \frac{g_2}{2} \Phi^* \cdot \Phi^* \hat{\varphi} \cdot \hat{\varphi} + H.c.$ is the

quadratic molecular Hamiltonian density, with $\tilde{\varepsilon}_\sigma = \hat{\varepsilon}_\sigma + g_{am}|\Phi|^2$ and $\tilde{\omega}_{ij} = (\tilde{\omega} + g_1|\Phi|^2)\delta_{ij} + g_1\Phi_j^*\Phi_i + 2g_2\Phi_i^*\Phi_j$.

To this quadratic order, the molecular and atomic excitations decouple and can therefore be diagonalized independently. The molecular part has been extensively studied in the context of spinor $F = 1$ condensates[28]. For the polar ($g_2 < 0$) MSF_p state, there are three “sound” modes, one Bogoliubov type with sound velocity $c_{||}^{(\text{MSF}_p)} = \sqrt{(g_1 + g_2)n_m/2m}$, and other two degenerate spin-waves with velocity $c_{\perp}^{(\text{MSF}_p)} = \sqrt{|g_2|n_m/2m}$. For the ferromagnetic ($g_2 > 0$) MSF_{fm} state, there is one Bogoliubov mode, with sound velocity $c^{(\text{MSF}_{fm})} = \sqrt{g_1 n_m/2m}$, one quadratic $k^2/2m$ ferromagnetic spin-wave mode, and one gapped quadratic $k^2/2m + 2g_2 n_m$ canonically conjugate mode.

The atomic sector can also be readily diagonalized, giving

$$E_{(a)}^{(\text{MSF})}(k) = \sqrt{\tilde{\varepsilon}_k^2 - \alpha^2 |\Phi \cdot \mathbf{k}|^2}, \quad (7)$$

where $\tilde{\varepsilon}_k = k^2/2m - \mu + g_{am}n_m$ and n_m is the molecular condensate density. The details of the spectrum only differ quantitatively between the MSF_p and MSF_{fm} phases, both exhibiting a minimum at a *finite* $\mathbf{k}_{\min}$ for $\nu > \nu_*$, and an atomic gap $\Delta(\nu) \equiv E_{(a)}^{(\text{MSF})}(k_{\min})$ that closes at the transition ν_{c1} into the corresponding AMSF states. Such spectrum (displaying a finite *k* minimum) should be experimentally observable through, e.g., rf spectroscopy and is also a complementary way to detect the approaching finite *Q* instability (phase transition) into the AMSF state. Simple analysis inside MSF_p gives

$$\begin{aligned} \mathbf{k}_{\min}^{(\text{MSF}_p)} &= \hat{n} \sqrt{(2m^2\alpha^2 + mg)n_m + m\nu}, \\ \Delta^{(\text{MSF}_p)} &= \sqrt{-(m\alpha^2 + g)m\alpha^2 n_m^2 - m\alpha^2 \nu n_m}, \\ \nu_*^{(\text{MSF}_p)} &= -(2m\alpha^2 + g)n_m, \\ \nu_{c1}^{(\text{MSF}_p - \text{AMSF}_p)} &= -(m\alpha^2 + g)n_m, \end{aligned} \quad (8)$$

with $g = g_1 + g_2 - 2g_{am}$ and we used lowest order MSF_p relation $\mu_m \equiv 2\mu - \nu \approx (g_1 + g_2)n_m$ to eliminate the atomic μ in favor of molecular condensate n_m and detuning ν . The corresponding expressions inside MSF_{fm} differ only slightly[23], except that $\mathbf{k}_{\min}^{(\text{MSF}_{fm})}$ lies in the $\hat{n}$ - $\hat{m}$ plane perpendicular to the ferromagnetic quantization axis, $\hat{\ell}$ rather than along it as in the MSF_p state.

Upon further increase of ν , the atomic gap $\Delta^{(\text{MSF})}(\nu)$ closes at ν_{c1} and atoms Bose-condense at a finite $\mathbf{k}_{\min}^{(\text{MSF})}(\nu_{c1})$ ($\mathbf{k}_{\min}^{(\text{MSF}_p)} = m\alpha\sqrt{n_m}$, $\mathbf{k}_{\min}^{(\text{MSF}_{fm})} = m\alpha\sqrt{n_m}/\sqrt{2}$), thereby breaking the remaining $Z_2 \times U_{\Delta N}(1)$ symmetry, with Z_2 corresponding to the discrete part of (atom number) $U_N(1)$ unbroken in the paired MSF states. The two associated order parameters are given by $\Psi_{\pm}(\mathbf{r}) = \sum_{\mathbf{Q}_n} (\pm \Psi_{\mathbf{Q}_n,1} e^{i(\mathbf{Q}_n \cdot \mathbf{r} - \theta_{\mathbf{Q}})} + \Psi_{-\mathbf{Q}_n,2}^* e^{i\mathbf{Q}_n \cdot \mathbf{r}})$, where $\theta_{\mathbf{Q}}$ is the phase of $\Delta_{\mathbf{Q}} \equiv \alpha \Phi \cdot \mathbf{Q} = |\alpha \Phi \cdot \mathbf{Q}| e^{i\theta_{\mathbf{Q}}}$ and $\mathbf{Q}_n = n\mathbf{Q}$

($n \in \mathcal{Z}$, an integer). The critical transition point for Ψ_+ and Ψ_- is split by $\pm|\Delta_{\mathbf{Q}}|$, respectively, so only Ψ_- condenses at ν_{c1} , allowing for the possibility of two transitions, MSF-AMSF $_-$ followed by AMSF $_-$ -AMSF $_+$. The values of order parameters and transition points can be straightforwardly worked out. Within MFT, we find that a single- Q state is energetically preferred[23], but we do not expect this to survive generically.

In addition to the molecular and atomic superfluidity, AMSF $_{+,-}^{p,fm}$ phases are finite-momentum condensates that generically can exhibit crystalline order and therefore are supersolids (at least in the case of more than one $\mathbf{Q}_n$ condensation[34]). For a collinear set of $\mathbf{Q}_n$'s, the supersolid is a unidirectional density wave breaking translational invariance along $\mathbf{Q}_n$, with the latter aligned with the quantization axis, $\hat{\mathbf{n}}$ of the MSF $_p$ state and transverse to the quantization axis $\hat{\ell}$ (i.e., lying in the $\hat{\mathbf{n}}-\hat{\mathbf{m}}$ plane) of the MSF $_{fm}$ state. Thus, while rotational $O(2)$ symmetry about the $\hat{\mathbf{n}}$ axis remains intact inside the AMSF $_p$ state, it is spontaneously broken by such a uniaxial density wave inside the AMSF $_{fm}$ state. It is notable that in this latter case, in the presence of fluctuations, such superfluid density wave will exhibit quantum liquid-crystal phenomenology similar to that of the fermion-paired Larkin-Ovchinnikov superfluid[23, 35]. The excitation spectra inside AMSF phases can be computed via a generalized Bogoliubov transformation and involve a diagonalization of a 10×10 matrix (corresponding to three and one coupled complex molecular and atomic fields), leading to Goldstone modes consistent with above symmetry-based arguments[23].

The nature of MSF $_{p,fm}$ -AMSF $_{p,fm}$ transitions (beyond MFT) remains an open question. Based on the experience with spinor condensates[32] and LO superfluid[35], we expect this system to exhibit a variety of fractional composite topological defects. We leave detailed study of these to a future publication[23].

Upon further increase of ν , the AMSF transitions into the ASF at $\nu_{c2} \approx (2\lambda - g_{am})n_a$ (with $\lambda = \frac{1}{4}(\lambda_1 + \lambda_2 + 2\lambda_{12})$ and the atomic condensate density n_a), determined by the point of vanishing of the $n_m(\nu)$ and $Q(\nu)$. The AMSF can thus exhibit a broad stability range $\nu_{c2} - \nu_{c1}$, set by a combination of $n_a, n_m, \lambda_i, g_i, \dots$, that in 3d we expect to survive beyond our mean-field analysis.

To summarize, we studied a degenerate gas of two-species bosonic atoms interacting through a p -wave Feshbach resonance, as realized in a ^{85}Rb - ^{87}Rb mixture. We showed that at intermediate FR detuning, such gas exhibits an atomic-molecular superfluid (AMSF) state condensed at a finite momentum, that undergoes phase transitions into a molecular p -wave (orbital spinor) superfluid (MSF) and into an s -wave atomic superfluid (ASF) at large negative and positive detunings, respectively. A magnetic field can be used to tune the modulation wave vector of the AMSF between zero and a value set by interactions as well as to drive quantum phase transitions

in this rich system.

We thank V. Gurarie for discussions and acknowledge support by the NSF No. DMR-0321848 (L. R., S. C.), Berkeley Miller, and University of Colorado Faculty Fellowships (L. R.). L. R. thanks Berkeley Physics Department for its hospitality during part of this work.

-
- [1] C. A. Regal *et al.*, Phys. Rev. Lett. **92**, 040403 (2004).
 - [2] M. W. Zwierlein *et al.*, Phys. Rev. Lett. **92**, 120403 (2004).
 - [3] J. Kinast *et al.*, Phys. Rev. Lett. **92**, 150402 (2004).
 - [4] C. Chin *et al.*, Science **305**, 1128 (2004).
 - [5] S. L. Cornish *et al.*, Phys. Rev. Lett. **85**, 1795 (2000).
 - [6] L. Radzihovsky *et al.*, Phys. Rev. Lett. **92**, 160402 (2004).
 - [7] L. Radzihovsky *et al.*, Ann. Phys. (N. Y.) **323**, 2376 (2008).
 - [8] M. W. J. Romans *et al.*, Phys. Rev. Lett. **93**, 020405 (2004).
 - [9] T.-L. Ho and R. B. Diener, Phys. Rev. Lett. **94**, 090402 (2005).
 - [10] V. Gurarie *et al.*, Phys. Rev. Lett. **94**, 230403 (2005).
 - [11] V. Gurarie and L. Radzihovsky, Ann. Phys. (N. Y.) **322**, 2 (2007).
 - [12] C. Cheng and S.-K. Yip, Phys. Rev. Lett. **95**, 070404 (2005).
 - [13] J. P. Gaebler *et al.*, Phys. Rev. Lett. **98**, 200403 (2007).
 - [14] J. Zhang *et al.*, Phys. Rev. A **70**, 030702 (2004).
 - [15] S. B. Papp *et al.*, Phys. Rev. Lett. **101**, 040402 (2008).
 - [16] P. Fulde and R. A. Ferrell, Phys. Rev. **135**, A550 (1964).
 - [17] A. I. Larkin and Y. N. Ovchinnikov, Sov. Phys. JETP **20**, 762 (1965).
 - [18] T. Mizushima *et al.*, Phys. Rev. Lett. **94** 060404 (2005).
 - [19] D. Sheehy and L. Radzihovsky, Phys. Rev. Lett. **96**, 060401 (2006).
 - [20] D. Sheehy and L. Radzihovsky, Ann. Phys. (N. Y.) **322**, 1790 (2007).
 - [21] For unequal masses FR interaction is given by $H_{FR} = \alpha \hat{\phi}^\dagger \cdot \left[\frac{m_1}{m_1+m_2} \hat{\psi}_1 (-i\nabla) \hat{\psi}_2 - \frac{m_2}{m_1+m_2} \hat{\psi}_2 (-i\nabla) \hat{\psi}_1 \right] + H.c.$ as required to preserve Galilean invariance. L.R. thanks V. Gurarie for discussion on this point.
 - [22] C. A. Regal *et al.*, Phys. Rev. Lett. **90**, 053201 (2003).
 - [23] S. Choi and L. Radzihovsky (to be published).
 - [24] B. D. Esry *et al.*, Phys. Rev. Lett. **78**, 3594 (1997).
 - [25] K.-S. Liu and M. E. Fisher, J. Low Temp. Phys. **10**, 655 (1973).
 - [26] J. M. Kosterlitz *et al.*, Phys. Rev. B **13**, 412 (1976).
 - [27] P. Calabrese *et al.*, Phys. Rev. B **67**, 054505 (2003).
 - [28] T.-L. Ho, Phys. Rev. Lett. **81**, 742 (1998).
 - [29] T. Ohmi and K. Machida, J. Phys. Soc. Jpn. **67**, 1822 (1998).
 - [30] D. M. Stamper-Kurn *et al.*, Phys. Rev. Lett. **80**, 2027 (1998).
 - [31] M. R. Matthews *et al.*, Phys. Rev. Lett. **81**, 243 (1998).
 - [32] S. Mukerjee, C. Xu, and J. E. Moore, Phys. Rev. Lett. **97**, 120406 (2006).
 - [33] $U_N(1)$ and $U_{\Delta N}(1)$ symmetries are associated with total atom number $N = N_1 + N_2 + 2N_m$ and difference $\Delta N = N_1 - N_2$ conservations, respectively.

- [34] In the single $\mathbf{Q}$ condensation (Fulde-Ferrell type[16]), the state (e.g., atom density) is uniform and the phase θ of Ψ_- is nothing but the superfluid phase. For the state with at least two $\pm\mathbf{Q}$ condensation (Larkin-Ovchinnikov type[17]), the two phases, $\theta_{\pm\mathbf{Q}}$ determine the superfluid phase $\theta = (\theta_{\mathbf{Q}} + \theta_{-\mathbf{Q}})/2$ and the density wave phonon $u = \frac{1}{2}(\theta_{\mathbf{Q}} - \theta_{-\mathbf{Q}})/Q$ [35].
- [35] L. Radzihovsky and A. Vishwanath, Phys. Rev. Lett. **103**, 010404 (2009).